


DESCRIPTIONS OF THE SOIL GROUPS

- 211** Silty clay loams to silt loams with (a) permeabilities from 1.0 to 2.0 inches per hour, (b) nearly level to very gentle slopes (maximum slopes 1 to 3 percent), and (c) depths to seasonal high water table less than 6 feet. These soils occur on low terraces and flood plains and are represented by the Zook-Leshara-Wann and Kennebec-Nodaway-Zook associations.
- 212** Silty clay loams to silt loams with (a) permeabilities from 1.0 to 2.0 inches per hour, (b) nearly level to very gentle slopes (maximum slopes 1 to 3 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are present on well-drained terraces and uplands in the central part of the State and are represented by the Hord-Hall and Holdrege-Hall associations.
- 222** Silty clay loams to silt loams with (a) permeabilities from 1.0 to 2.0 inches per hour, (b) nearly level to strong slopes (maximum slopes 3 to 10 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are widely distributed throughout the State and are represented by the Holdrege and Keith-Alliance-Rosebud associations.
- 232** Silty clay loams to loams with (a) permeabilities from 1.0 to 2.0 inches per hour, (b) very gentle to moderately steep slopes (maximum slopes 10 to 20 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are widely distributed and significant within the State and are represented by the Holdrege-Coly-Uly and Ulysses-Keith-Colby associations.
- 252** Silty clay loams to loams with (a) permeabilities from 1.0 to 2.0 inches per hour, (b) gentle to very steep slopes (maximum slopes that exceed 30 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are common in central and southwestern portions of the State and are represented by the Colby-Ulysses and Colby-Uly associations.
- 311** Silt loams to fine sandy loams with (a) permeabilities from 2.0 to 5.0 inches per hour, (b) nearly level to very gentle slopes (maximum slopes 1 to 3 percent), and (c) depths to seasonal high water table less than 6 feet. These soils are present in the Platte and Missouri River flood plains and are represented by the Gibbon-Wann and Lawet-Wann-Lex associations.
- 322** Silt loams to fine sandy loams with (a) permeabilities from 2.0 to 5.0 inches per hour, (b) nearly level to strong slopes (maximum slopes 3 to 10 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are common in transitional areas between the sandhills and silty uplands and are represented by the Moody-Bazile-Trent and Jayem-Haxton-Rosebud associations.
- 411** Loams to fine sands with (a) permeabilities from 5.0 to 10.0 inches per hour, (b) nearly level to very gentle slopes (maximum slopes 1 to 3 percent), and (c) depths to seasonal high water table less than 6 feet. These soils are along flood plains and are represented by the Las-Las Animas-McCook and Lawet-Elsmere-Gannett associations.
- 422** Fine sandy loams to fine sands with (a) permeabilities from 5.0 to 10 inches per hour, (b) nearly level to strong slopes (maximum slopes 3 to 10 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils occur on uplands, terraces, and footslopes in transitional areas between sandy and silty soils and are represented by the Bazile-Paka-Thurman and Jayem-Sarben-Valent associations.
- 511** Fine sandy loams to fine sands with (a) permeabilities exceeding 10.0 inches per hour, (b) nearly level to very gentle slopes (maximum slopes 1 to 3 percent), and (c) shallow water tables with depths to seasonal high water table less than 6 feet. These soils are on flood plains and in Sand Hills valleys and are represented by the Gothenburg-Platte and Loup-Elsmere-Dunday associations.
- 532** Loamy fine sands to fine sands with (a) permeabilities exceeding 10.0 inches per hour, (b) nearly level to steep slopes (maximum slopes 10 to 20 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are represented by the Jansen-Meadin and Valentine-Hersh associations.
- 541** Loamy fine sands to fine sands with (a) permeabilities exceeding 10.0 inches per hour, (b) nearly level to very steep slopes (maximum slopes 20 to 30 percent), and (c) substantial areas having depths to seasonal high water table less than 6 feet. This hydrologic soil group is rather unique in that steeply sloping dunes alternate with subirrigated valleys with shallow water tables and seasonal ponding. The Valentine-Els and Valentine-Elsmere-Gannett associations represent these soils.
- 542** Loamy fine sands to fine sands with (a) permeabilities exceeding 10.0 inches per hour, (b) nearly level to very steep slopes (maximum slopes 20 to 30 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are principally found in the Sand Hills uplands and are represented by the Valentine-Tassel and Valentine-Simon associations.
- 552** Loamy fine sands to fine sands with (a) permeabilities exceeding 10.0 inches per hour, (b) gentle to very steep slopes (maximum slopes exceeding 30 percent), and (c) depths to seasonal high water table exceeding 6 feet. These soils are the most prevalent upland soils of the Sand Hills and are represented by the Valentine and Valentine, hilly and rolling associations.

HYDROLOGIC CHARACTERISTICS OF THE SOIL GROUPS

Soil group	Average permeability of 60-inch soil profile (inches per hour)	Average permeability of least permeable horizon (inches per hour)	Average available water capacity (inches per inch)	Average maximum soil slope (percent)	Depth to seasonal high water table (feet)
211	1.52	.73	.19	2	<6
212	1.23	1.14	.20	3	>6
222	1.23	1.09	.20	5	>6
232	1.28	1.21	.19	15	>6
252	1.31	1.27	.21	48	>6
311	3.61	.93	.16	2	<6
322	2.91	2.02	.16	3	>6
411	8.42	1.52	.13	2	<6
422	6.85	4.15	.13	8	>6
511	12.90	3.99	.09	3	<6
532	12.46	5.21	.09	14	>6
541	12.67	11.38	.08	27	<6
542	12.20	7.57	.08	27	>6
552	12.38	10.67	.07	50	>6



HYDROLOGIC SOIL GROUPS IN THE
NORTH PLATTE QUADRANGLE, NEBRASKA